



REVISION NOTES · CLASS 11 AND 12

NEET Biology High-Yield Notes

NEET biology rewards exact NCERT facts. These notes collect the numbers, names and ratios that turn up again and again, chapter by chapter, so you can revise the whole syllabus in an evening.

NEET

CBSE 12

01 The cell

Cell theory – Schleiden and Schwann; Virchow added that new cells come from existing cells (omnis cellula e cellula)

Ribosomes – prokaryotes 70S (50S + 30S); eukaryotes 80S (60S + 40S)

Semi-autonomous organelles – mitochondria and chloroplasts have their own circular DNA and 70S ribosomes

Membrane model – fluid mosaic model, Singer and Nicolson (1972)

Centrioles and cilia – centriole 9 + 0 triplets; cilia and flagella 9 + 2 axoneme

Lysosomes – hydrolytic enzymes, active at acidic pH

02 Cell cycle and division

Interphase – G₁, S (DNA replicates), G₂

Mitosis – metaphase is the best stage to study chromosome morphology

Prophase I of meiosis – synapsis in zygotene; crossing over in pachytene; chiasmata seen in diplotene; terminalisation in diakinesis

Meiosis I – reductional division; meiosis II is equational

03 Molecular biology

DNA structure – Watson and Crick (1953); B-DNA has about 10 bp per turn and a pitch of 3.4 nm

Base pairing – A = T (2 hydrogen bonds); G ≡ C (3 hydrogen bonds); Chargaff's rule A + G = T + C

Replication – semi-conservative (Meselson and Stahl, using ¹⁵N); Okazaki fragments form on the lagging strand

Genetic code – triplet, degenerate, nearly universal; start codon AUG; stop codons UAA, UAG, UGA

Lac operon – Jacob and Monod; lactose (allolactose) is the inducer

Human genome (NCERT figures) – 3164.7 million bp; about 30,000 genes; under 2 per cent codes for proteins

04 Genetics

Mendelian ratios – monohybrid F₂ 3:1 (genotype 1:2:1); dihybrid 9:3:3:1; test cross 1:1

Incomplete dominance – snapdragon flower colour, F₂ 1:2:1

Co-dominance – ABO blood groups (I^A, I^B, i)

Sex determination – humans XX-XY; grasshopper XX-XO; birds ZW female, ZZ male

X-linked recessive – haemophilia, colour blindness

Sickle-cell anaemia – glutamic acid replaced by valine at position 6 of β-globin

Chromosomal disorders – Down syndrome trisomy 21; Klinefelter 47,XXY; Turner 45,XO

05 Human physiology

Muscle contraction – sliding filament theory; Ca²⁺ binds troponin, exposing actin's binding sites to myosin heads; ATP powers the cross-bridges

Nerve impulse – the Na⁺/K⁺ pump moves 3 Na⁺ out for every 2 K⁺ in; chemical synapses pass the signal with neurotransmitters such as acetylcholine

Breathing – tidal volume about 500 mL; vital capacity = IRV + TV + ERV

Gas transport – O₂ about 97% as oxyhaemoglobin; CO₂ about 70% as bicarbonate, 20–25% as carbaminohaemoglobin, 7% dissolved

Heart – SA node is the pacemaker (70–75 beats/min); cardiac output ≈ 70 mL × 72 ≈ 5 L/min; normal BP 120/80 mm Hg

ECG – P wave atrial depolarisation; QRS ventricular depolarisation; T wave ventricular repolarisation

Kidney – GFR about 125 mL/min (180 L/day); PCT reabsorbs 70–80% of water and electrolytes; ADH acts on DCT and collecting duct

Hormones – insulin (β cells) lowers blood glucose, glucagon (α cells) raises it; ADH and oxytocin are made in the hypothalamus and released by the posterior pituitary

06 Plant physiology

Photosystems – PS I reaction centre P700; PS II P680; water splits at PS II

C₃ and C₄ – Calvin cycle's first stable product is 3-PGA; Hatch-Slack (C₄) first product is OAA, with Kranz anatomy (maize, sugarcane)

RuBisCO – the most abundant enzyme on Earth; its oxygenase activity causes photorespiration in C₃ plants

Respiration – glycolysis gives a net 2 ATP; Krebs cycle in the matrix; ETS on the inner membrane; NCERT net gain 38 ATP per glucose

Respiratory quotient – carbohydrates 1; fats about 0.7; proteins about 0.9

Plant hormones – auxin (apical dominance, rooting); gibberellins (bolting, malting); cytokinins (cell division, delay ageing); ethylene (ripening); ABA (stress, stomatal closure, dormancy)

07 Reproduction

Flowering plants – double fertilisation, syngamy plus triple fusion (triploid endosperm)

Menstrual cycle – about 28 days; ovulation about day 14 after the LH surge; corpus luteum secretes progesterone

Testes – Sertoli cells nourish developing sperm; Leydig cells secrete testosterone

Pregnancy – implantation at the blastocyst stage; hCG maintains the corpus luteum

08 Ecology

Energy flow – Lindeman's 10 per cent law; the pyramid of energy is always upright

Inverted pyramids – biomass in the sea; numbers in a single tree

Species-area relationship – $\log S = \log C + Z \log A$, Z usually 0.1 to 0.2

Biodiversity – 34 hotspots worldwide; India has the Western Ghats-Sri Lanka, Indo-Burma and Himalaya

Rivet popper hypothesis – Paul Ehrlich

Causes of loss (the Evil Quartet) – habitat loss and fragmentation, over-exploitation, alien species invasions, co-extinctions

Conservation – in situ (biosphere reserves, national parks, sanctuaries, sacred groves); ex situ (zoos, botanical gardens, seed banks, cryopreservation)

09 Biotechnology and health

Restriction enzyme – EcoRI cuts at GAATTC

PCR – denaturation, annealing, extension with Taq polymerase (*Thermus aquaticus*)

Vector pBR322 – ori, ampicillin and tetracycline resistance genes

Bt crops – cry genes from *Bacillus thuringiensis*; cryI_{Ac} and cryII_{Ab} for cotton, cryI_{Ab} for corn

RNA interference – protects tobacco from the nematode *Meloidogyne incognita*

Firsts – gene therapy for ADA deficiency (1990); Rosie, the transgenic cow whose milk carried human α -lactalbumin (1997)

Diseases – malaria, *Plasmodium* via female Anopheles; typhoid, *Salmonella typhi* (Widal test); dengue and chikungunya via Aedes; filariasis, *Wuchereria* via Culex

AIDS – caused by HIV, a retrovirus that destroys helper T cells; ELISA detects infection

How to use these notes

- 1 Read the matching NCERT paragraph after each section: NEET often lifts the exact sentence.
- 2 Make your own list of examples (organisms, diseases, hormones); they are easy marks.
- 3 Revise diagrams with labels: they carry both one-mark and assertion-reason questions.

ACHIEVERS · SINCE 1999

Want a Biology tutor for NEET?

One-to-one tutors at home in the Chandigarh Tricity, or live online anywhere. Our office confirms the tutor and the fee before the first class.

WhatsApp +91 86990 09555
hometutorsindia.com



Scan to WhatsApp



REVISION NOTES · CLASS 11 AND 12

Chemistry Quick Revision for JEE and NEET

Physical chemistry formulas, inorganic trends and the organic reactions examiners return to most often, condensed onto a few pages for the last weeks before JEE or NEET.

JEE

NEET

CBSE 12

01 Mole concept and solutions

Moles – $n = \text{mass/molar mass} = \text{number of particles}/N_A$

Molarity – $M = \text{moles of solute/litres of solution}$; **molality** $m = \text{moles of solute/kg of solvent}$

Mole fraction – $x_A = n_A/(n_A + n_B)$

Normality – $N = M \times n\text{-factor}$; **dilution** $M_1V_1 = M_2V_2$

Raoult's law – $p_A = x_A p_A^\circ$; **relative lowering** $(p^\circ - p)/p^\circ = x_{\text{solute}}$

Colligative properties – $\Delta T_b = iK_b m$; $\Delta T_f = iK_f m$; **osmotic pressure** $\pi = iCRT$

Henry's law – $p = K_H x$

02 Atomic structure

Bohr energy – $E_n = -13.6 Z^2/n^2 \text{ eV}$; **radius** $r_n = 0.529 n^2/Z \text{ \AA}$

de Broglie – $\lambda = h/mv$; **Heisenberg** $\Delta x \cdot \Delta p \geq h/4\pi$

Quantum numbers – n ; l from 0 to $n - 1$; m_l from $-l$ to $+l$; $m_s = \pm 1/2$

Capacity – $2n^2$ electrons per shell; $2l + 1$ orbitals per subshell

Nodes – radial $n - l - 1$; angular l

Filling order – $(n + l)$ rule, then Pauli exclusion and Hund's rule

03 Gases

Ideal gas – $PV = nRT$, $R = 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1} = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$

Dalton – $p_i = x_i p_{\text{total}}$

Graham – rate $\propto 1/\sqrt{M}$

van der Waals – $(P + an^2/V^2)(V - nb) = nRT$

Compressibility – $Z = PV/nRT$

Molecular speeds – $u_{\text{rms}} = \sqrt{3RT/M}$, $u_{\text{avg}} = \sqrt{8RT/\pi M}$, $u_{\text{mp}} = \sqrt{2RT/M}$

04 Thermodynamics

First law – $\Delta U = q + w$ ($w = \text{work done on the system}$); $w = -P_{\text{ext}}\Delta V$

Enthalpy – $\Delta H = \Delta U + \Delta n_g RT$; **Hess's law** adds reaction enthalpies

Gibbs energy – $\Delta G = \Delta H - T\Delta S$; spontaneous when $\Delta G < 0$

Link to equilibrium – $\Delta G^\circ = -RT \ln K = -2.303RT \log K$

Link to cells – $\Delta G^\circ = -nFE^\circ_{\text{cell}}$

05 Equilibrium

K_p and K_c – $K_p = K_c(RT)^{\Delta n}$

pH – $\text{pH} = -\log[H^+]$; $\text{pH} + \text{pOH} = 14$ and $K_w = 10^{-14}$ at 25°C

Weak acid – $[H^+] = \sqrt{K_a C}$, **degree of dissociation** $\alpha = \sqrt{K_a/C}$

Buffer (Henderson) – $\text{pH} = \text{p}K_a + \log([\text{salt}]/[\text{acid}])$

Salt of weak acid and strong base – $\text{pH} = 7 + \frac{1}{2}(\text{p}K_a + \log C)$

Solubility product – $K_{\text{sp}} = s^2$ for AB ; $4s^3$ for AB_2

06 Electrochemistry and kinetics

Cell emf – $E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$

Nernst equation (298 K) – $E = E^\circ - (0.0591/n) \log Q$

Faraday's law – $m = (M/nF)It$, $F = 96,500 \text{ C mol}^{-1}$

Conductivity – $\Lambda_m = \kappa \times 1000/M$ (κ in S cm^{-1}); **Kohlrausch** $\Lambda_m^\circ = v_+ \lambda_+^\circ + v_- \lambda_-^\circ$

Zero order – $t_{1/2} = [A]_0/2k$

First order – $k = (2.303/t) \log([A]_0/[A])$, $t_{1/2} = 0.693/k$

Arrhenius – $k = Ae^{-E_a/RT}$; $\log(k_2/k_1) = [E_a/2.303R](1/T_1 - 1/T_2)$

07 Periodic trends and bonding

Atomic radius – decreases across a period, increases down a group

Ionisation enthalpy – increases across a period; exceptions $N > O$, $Be > B$, $Mg > Al$

Electronegativity – fluorine is highest (4.0 on the Pauling scale); chlorine has the most negative electron gain enthalpy

Bond order (MOT) – $(N_b - N_a)/2$; O_2 has bond order 2 and is paramagnetic; N_2 has bond order 3

Hybridisation – 2 sp linear; 3 sp² trigonal planar; 4 sp³ tetrahedral; 5 sp³d trigonal bipyramidal; 6 sp³d² octahedral

Shapes – H_2O bent 104.5°; NH_3 pyramidal 107°; SF_4 see-saw; ClF_3 T-shaped; XeF_2 linear; XeF_4 square planar

Formal charge – valence electrons – non-bonding electrons – $\frac{1}{2}$ bonding electrons

08 Coordination compounds

Crystal field (octahedral) – $t_{2g} -0.4\Delta_o$, $e_g +0.6\Delta_o$

Spectrochemical series – $I^- < Br^- < Cl^- < F^- < OH^- < H_2O < NH_3 < en < CN^- < CO$

Magnetic moment (spin only) – $\mu = \sqrt{n(n+2)}$ BM, n = unpaired electrons

09 Named reactions

Aldol – aldehydes or ketones with α -H in dilute alkali give β -hydroxy carbonyl compounds

Cannizzaro – aldehydes without α -H in concentrated alkali give an alcohol and a carboxylate

Wurtz – $2R-X + 2Na$ in dry ether give R-R

Friedel-Crafts – alkylation or acylation of benzene with $AlCl_3$

Reimer-Tiemann – phenol + $CHCl_3$ + NaOH give salicylaldehyde

Kolbe – sodium phenoxide + CO_2 give salicylic acid

Hell-Volhard-Zelinsky – carboxylic acid + X_2 /red P gives the α -halo acid

Hoffmann bromamide – amide + $Br_2/NaOH$ gives an amine with one carbon fewer

Sandmeyer – diazonium salt + CuX gives an aryl halide

Gabriel phthalimide – a route to pure primary amines

Clemmensen and Wolff-Kishner – reduce $C=O$ to CH_2 ($Zn-Hg/HCl$; NH_2NH_2/KOH)

Rosenmund – acid chloride + $H_2/Pd-BaSO_4$ gives an aldehyde

Etard – toluene + CrO_2Cl_2 gives benzaldehyde

Williamson – $RO^-Na^+ + R'X$ gives the ether $R-O-R'$

10 Organic rules of thumb

Markovnikov – H adds to the carbon with more hydrogens; HBr with peroxide adds anti-Markovnikov

Substitution – S_N1 favours $3^\circ > 2^\circ > 1^\circ$ and racemises; S_N2 favours $1^\circ > 2^\circ > 3^\circ$ with inversion

Carbocation stability – $3^\circ > 2^\circ > 1^\circ > \text{methyl}$

Acidity – carboxylic acids > phenols > water > alcohols; electron-withdrawing groups raise acidity

11 Laboratory tests

Tollens' reagent – silver mirror with aldehydes

Fehling's solution – red precipitate with aliphatic aldehydes

Iodoform test – CH_3CO- or $CH_3CH(OH)-$ groups give yellow CHI_3

Lucas test – tertiary alcohols turn cloudy at once

Carbylamine test – primary amines give a foul-smelling isocyanide

Neutral $FeCl_3$ – violet colour with phenols

2,4-DNP – orange precipitate with aldehydes and ketones

Hinsberg's reagent – tells primary, secondary and tertiary amines apart

How to use these notes

- 1 Learn named reactions as reagent $\rightarrow$ product pairs, then test yourself by covering one side.
- 2 For trends, remember the exceptions: they are what the questions ask about.
- 3 NCERT lines are the source for most inorganic questions; read the relevant chapter after this sheet.

ACHIEVERS · SINCE 1999

Want a Chemistry tutor for JEE or NEET?

One-to-one tutors at home in the Chandigarh Tricity, or live online anywhere. Our office confirms the tutor and the fee before the first class.

WhatsApp +91 86990 09555
hometutorsindia.com



Scan to WhatsApp



FORMULA SHEET · CLASS 11 AND 12

Physics Formula Sheet for JEE and NEET

The formulas that carry most of the marks in JEE Main, JEE Advanced and NEET physics, grouped the way the NCERT books teach them. Print it, stick it above your desk, and test yourself on one section a day.

JEE

NEET

CBSE 12

01 Units and errors

Force – $[M L T^{-2}]$

Work, energy – $[M L^2 T^{-2}]$

Power – $[M L^2 T^{-3}]$

Pressure, stress – $[M L^{-1} T^{-2}]$

Gravitational constant G – $[M^{-1} L^3 T^{-2}]$

Planck's constant h – $[M L^2 T^{-1}]$

Error in $x = a^p b^q / c^r$ – $\Delta x/x = p(\Delta a/a) + q(\Delta b/b) + r(\Delta c/c)$

02 Kinematics

Equations of motion – $v = u + at$, $s = ut + \frac{1}{2}at^2$, $v^2 = u^2 + 2as$

Distance in the n th second – $s_n = u + a(n - \frac{1}{2})$

Projectile – $T = 2u \sin\theta/g$, $H = u^2 \sin^2\theta/2g$, $R = u^2 \sin 2\theta/g$ (maximum at 45°)

Relative velocity – $v_{AB} = v_A - v_B$

Uniform circular motion – $a_c = v^2/r = \omega^2 r$, $v = \omega r$

03 Laws of motion and friction

Newton's second law – $F = dp/dt = ma$

Impulse – $J = F\Delta t = \Delta p$

Friction – $f_s \leq \mu_s N$, $f_k = \mu_k N$, angle of repose: $\tan\theta = \mu_s$

Banked road (no friction) – $\tan\theta = v^2/rg$

Level curve, maximum speed – $v = \sqrt{\mu rg}$

04 Work, energy and power

Work – $W = F \cdot s = Fs \cos\theta$

Kinetic energy – $K = \frac{1}{2}mv^2 = p^2/2m$

Potential energy – mgh (near Earth), $\frac{1}{2}kx^2$ (spring)

Work-energy theorem – $W_{net} = \Delta K$

Power – $P = dW/dt = F \cdot v$

Elastic collision (1D) – $v_1 = [(m_1 - m_2)u_1 + 2m_2u_2]/(m_1 + m_2)$

Coefficient of restitution – $e = (v_2 - v_1)/(u_1 - u_2)$

05 Rotational motion

Torque – $\tau = r \times F = I\alpha$

Angular momentum – $L = I\omega = r \times p$

Rotational kinetic energy – $\frac{1}{2}I\omega^2$

Parallel axis – $I = I_{cm} + Md^2$

Perpendicular axis (lamina) – $I_z = I_x + I_y$

Moments of inertia – ring MR^2 ; disc $\frac{1}{2}MR^2$; solid sphere $(2/5)MR^2$; hollow sphere $(2/3)MR^2$; rod about centre $ML^2/12$, about end $ML^2/3$

Rolling without slipping – $v = \omega R$, $K = \frac{1}{2}mv^2(1 + k^2/R^2)$

Rolling down an incline – $a = g \sin\theta/(1 + k^2/R^2)$

06 Gravitation

Newton's law – $F = Gm_1m_2/r^2$

Surface gravity – $g = GM/R^2$

At height h ($h \ll R$) – $g' \approx g(1 - 2h/R)$

At depth d – $g' = g(1 - d/R)$

Orbital speed – $v_o = \sqrt{GM/r}$

Escape speed – $v_e = \sqrt{2GM/R} = \sqrt{2gR} \approx 11.2$ km/s for Earth

Kepler's third law – $T^2 = 4\pi^2 r^3/GM$

Potential energy – $U = -GMm/r$

07 Solids and fluids

Young's modulus – $Y = (F/A)/(\Delta L/L)$

Pressure at depth – $P = P_0 + \rho gh$

Buoyant force – $F_B = \rho_{fluid} Vg$

Continuity – $A_1 v_1 = A_2 v_2$

Bernoulli – $P + \frac{1}{2} \rho v^2 + \rho gh = \text{constant}$

Stokes' law – $F = 6\pi\eta r v$

Terminal velocity – $v_t = 2r^2(\rho - \sigma)g/9\eta$

Excess pressure – drop $2T/r$, soap bubble $4T/r$

Capillary rise – $h = 2T \cos\theta/\rho g r$

08 Heat and thermodynamics

Thermal expansion – $\Delta L = \alpha L \Delta T$, $\beta = 2\alpha$, $\gamma = 3\alpha$

Heat – $Q = mc\Delta T$, $Q = mL$ (change of state)

First law – $\Delta U = Q - W$ (W = work done by the gas)

Isothermal work – $W = nRT \ln(V_2/V_1)$

Adiabatic – $PV^\gamma = \text{constant}$, $W = (P_1 V_1 - P_2 V_2)/(\gamma - 1)$

Specific heats – $C_p - C_v = R$, $\gamma = C_p/C_v$ (monatomic 5/3, diatomic 7/5)

Efficiency – $\eta = W/Q_1$; Carnot $\eta = 1 - T_2/T_1$

Refrigerator – $\text{COP} = T_2/(T_1 - T_2)$

Kinetic theory – $P = \frac{1}{3} \rho v_{\text{rms}}^2$, $v_{\text{rms}} = \sqrt{3RT/M}$, average KE per molecule = $(3/2)kT$

Radiation – Stefan $E = \sigma e T^4$; Wien $\lambda_m T = b \approx 2.9 \times 10^{-3} \text{ m K}$

Newton's law of cooling – $dT/dt = -k(T - T_0)$

09 Oscillations and waves

SHM – $x = A \sin(\omega t + \phi)$, $v = \omega \sqrt{A^2 - x^2}$, $a = -\omega^2 x$

Time period – spring $T = 2\pi\sqrt{m/k}$; simple pendulum $T = 2\pi\sqrt{L/g}$

Energy in SHM – $E = \frac{1}{2} k A^2$

Wave speed – $v = f\lambda$; on a string $v = \sqrt{T/\mu}$; sound in a gas $v = \sqrt{\gamma RT/M}$

Open pipe or string – $f_n = nv/2L$

Closed pipe – $f_n = (2n - 1)v/4L$ (odd harmonics only)

Beats – $f_{\text{beat}} = |f_1 - f_2|$

Doppler effect – $f' = f(v \pm v_o)/(v \mp v_s)$ (upper signs: moving towards each other)

10 Electrostatics

Coulomb's law – $F = kq_1 q_2 / r^2$, $k = 1/4\pi\epsilon_0 \approx 9 \times 10^9 \text{ N m}^2 \text{ C}^{-2}$

Point charge – $E = kq/r^2$, $V = kq/r$

Dipole – axial $E = 2kp/r^3$, equatorial $E = kp/r^3$, torque $\tau = p \times E$

Gauss's law – $\Phi = q_{\text{enc}}/\epsilon_0$

Line charge – $E = \lambda/2\pi\epsilon_0 r$

Infinite sheet – $E = \sigma/2\epsilon_0$

Capacitance – $C = Q/V$; parallel plates $C = \epsilon_0 A/d$

Energy stored – $U = \frac{1}{2} CV^2 = Q^2/2C$

Combinations – series $1/C = \sum 1/C_i$; parallel $C = \sum C_i$

11 Current electricity

Current – $I = dq/dt = nAev_d$

Ohm's law – $V = IR$, $R = \rho L/A$

Temperature – $R_T = R_0(1 + \alpha\Delta T)$

Combinations – series $R = \sum R_i$; parallel $1/R = \sum 1/R_i$

Power – $P = VI = I^2 R = V^2/R$

Cell – $V = E - Ir$; maximum power to the load when $R = r$

Kirchhoff – $\sum I = 0$ at a junction; $\sum V = 0$ round a loop

Wheatstone bridge (balanced) – $P/Q = R/S$

12 Magnetism and induction

Force on a charge – $F = q(v \times B)$; radius $r = mv/qB$; period $T = 2\pi m/qB$

Force on a wire – $F = IL \times B$

Biot-Savart – $dB = (\mu_0/4\pi) I dl \times \hat{r}/r^2$

Long straight wire – $B = \mu_0 I/2\pi r$

Centre of a loop – $B = \mu_0 I/2R$

Solenoid – $B = \mu_0 nI$

Parallel wires, force per length – $\mu_0 I_1 I_2 / 2\pi d$

Faraday's law – $\epsilon = -d\Phi/dt$; motional emf $\epsilon = Blv$

Inductance – $U = \frac{1}{2} LI^2$; solenoid $L = \mu_0 n^2 A l$

Growth in an LR circuit – $I = I_0(1 - e^{-t/\tau})$, $\tau = L/R$; RC circuit $\tau = RC$

13 Alternating current

RMS value – $V_{\text{rms}} = V_0/\sqrt{2}$

Reactance – $X_L = \omega L$, $X_C = 1/\omega C$

Impedance – $Z = \sqrt{R^2 + (X_L - X_C)^2}$

Resonance – $f_0 = 1/(2\pi\sqrt{LC})$, $Q = \omega_0 L/R$

Power – $P = V_{\text{rms}} I_{\text{rms}} \cos\phi$

Ideal transformer – $V_s/V_p = N_s/N_p = I_p/I_s$

14 Optics

Mirror – $1/v + 1/u = 1/f$, $f = R/2$, $m = -v/u$

Lens – $1/v - 1/u = 1/f$, $m = v/u$, $P = 1/f$ (dioptries, f in metres)

Lens maker's formula – $1/f = (\mu - 1)(1/R_1 - 1/R_2)$

Lenses in contact – $1/F = 1/f_1 + 1/f_2$

Snell's law – $n_1 \sin i = n_2 \sin r$; critical angle $\sin C = 1/\mu$

Prism – $\mu = \sin[(A + \delta_m)/2]/\sin(A/2)$; thin prism $\delta = (\mu - 1)A$

Young's double slit – fringe width $\beta = \lambda D/d$

Single slit, first minimum – $a \sin \theta = \lambda$

Polarisation – Malus $I = I_0 \cos^2 \theta$; Brewster $\tan i_p = \mu$

Instruments – telescope $m = f_o/f_e$; compound microscope $m \approx (L/f_o)(D/f_e)$

15 Modern physics

Photon – $E = hf = hc/\lambda$, $hc \approx 1240 \text{ eV nm}$

Photoelectric effect – $K_{\max} = hf - \phi = eV_0$

de Broglie – $\lambda = h/p = h/\sqrt{2mK}$; electron through V volts: $\lambda \approx 1.227/\sqrt{V} \text{ nm}$

Bohr model – $r_n = 0.529 n^2/Z \text{ \AA}$, $E_n = -13.6 Z^2/n^2 \text{ eV}$

Spectral lines – $1/\lambda = RZ^2(1/n_1^2 - 1/n_2^2)$, $R \approx 1.097 \times 10^7 \text{ m}^{-1}$

How to use this sheet

- 1 Cover the right-hand side of each line and say the formula out loud before you look.
- 2 Write the units next to every answer: it catches half of all slips.
- 3 Pair each formula with one solved question from your NCERT book or a previous paper.

Radioactive decay – $N = N_0 e^{-\lambda t}$, $T_{1/2} = 0.693/\lambda$, mean life $\tau = 1/\lambda$

Mass-energy – $E = mc^2$; $1 \text{ u} = 931.5 \text{ MeV}$

Nuclear radius – $R = R_0 A^{1/3}$

Diodes – a p-n junction conducts in forward bias above its barrier (about 0.7 V for silicon); a Zener diode in reverse breakdown regulates voltage

Logic and rectifiers – NAND and NOR are universal gates; full-wave output frequency = $2 \times$ input

16 Constants worth knowing

$g = 9.8 \text{ m s}^{-2}$	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
$c = 3 \times 10^8 \text{ m s}^{-1}$	$h = 6.63 \times 10^{-34} \text{ J s}$
$e = 1.6 \times 10^{-19} \text{ C}$	$m_e = 9.1 \times 10^{-31} \text{ kg}$
$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$	$R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$
$k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$	$\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$
$\mu_0 = 4\pi \times 10^{-7} \text{ T m A}^{-1}$	$\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$
$1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$	

ACHIEVERS · SINCE 1999

Want a Physics tutor for JEE or NEET?

One-to-one tutors at home in the Chandigarh Tricity, or live online anywhere. Our office confirms the tutor and the fee before the first class.

WhatsApp +91 86990 09555
[hometutorsindia.com](https://www.hometutorsindia.com)



Scan to WhatsApp

Free to print and share for study. Please keep the Achievers name and number on every page. Exam patterns change, so always check the details with the official board or testing agency. © 2026 Achievers Home Tutions · <https://www.hometutorsindia.com/physics-formula-sheet-jee-neet.html>